

Control of *Alternaria alternata* Using Melaleuca Essential Oil (*Melaleuca alternifolia*)

**Flávia Mota de Figuerêdo Alves¹, Kevison Romulo da Silva França^{1*},
Ionaly Gomes de Araújo¹, Lídia Pinheiro da Nóbrega¹,
Alda Leaby dos Santos Xavier¹, Tiago Silva Lima²,
Ana Paula Medeiros dos Santos Rodrigues³,
Antônio Francisco de Mendonça Júnior⁴
and Tiago Augusto Lima Cardoso⁵**

¹Department of Agroindustrial Systems, Federal University of Campina Grande, Pombal, PB, Brazil.

²Department of Plant Protection, Federal University of Alagoas, Alagoas, AL, Brazil.

³Department of Agronomy and Plant Protection, Federal Rural University of Semiarid, Mossoró, RN, Brazil.

⁴Department of Agronomy, Rural Federal University of Pernambuco, Recife, PE, Brazil.

⁵Laboratory of Phytopathology, Federal University of Campina Grande, Pombal, PB, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. Authors FMFA and KRSF carried out the experiment, managed the analyses of the study, performed calculations, managed the literature searches, interpreted the results, wrote the first draft of the manuscript and finished the paper with the others authors suggestions. Authors IGA, LPN, ALSX and TSL helped in carrying out the experiment, with results interpretation, managing the analyses, discussion and writing the paper. Authors AFMJ and APMSR designed the study, helped in carrying out the experiment, performed calculations and did the review of full paper. Author TALC helped in designing the study, performed the statistical analysis, helped with the discussion, did the review of full paper and made the translation into English. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/v40i330364

Editor(s):

(1) Dr. Mohamed Fadel, Professor, Microbial Chemistry, Department of Genetic Engineering and Biotechnology, Division National Research Center, Egypt.

Reviewers:

(1) Jacek Łyczko, Wrocław University of Environmental and Life Sciences, Poland.

(2) Jamile da Silva Oliveira, Brazil.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50927>

Original Research Article

**Received 09 June 2019
Accepted 22 August 2019
Published 28 August 2019**

ABSTRACT

Aims: This study aimed to evaluate the fungitoxic potential of melaleuca essential oil on the mycelial growth of *Alternaria alternata* under *in vitro* condition and the treatment of cowpea beans.

Study Design: The experiments comprised completely randomized designs: Eleven treatments with five replicates on *in vitro* test; and six treatments with five replicates on *in vivo* test.

Place and Duration of Study: The work was carried out at the Center for Agrifood Science and Technology of the Federal University of Campina Grande, Pombal, Brazil, since February 2018 to February 2019.

Methodology: In the *in vitro* experiment, the essential oil was incorporated into the culture medium and poured into Petri dishes. The treatments consisted of different concentrations of the essential oil (0.0125, 0.025, 0.05, 0.1, 0.2, 0.25, 0.50, 0.75, and 1.0%), a negative control (0.0%), and a positive control (Thiram). Discs of culture medium with fungal mycelia were inoculated in the center of the plates and incubated for seven days at 27±2°C. The percentage of mycelial growth inhibition (PGI) and the index of mycelial growth speed (IMGS) was calculated to verify the difference between treatments. In the *in vivo* experiment, the bean seeds were treated with different concentrations of EO (0.0, 0.2, 0.5, 1.0, and 5.0%), a negative control (0.0%), and positive control (Thiram). Seeds were inoculated with colonies of the fungus for 48 hours, and after that, we performed the seed sanity test.

Results: Under *in vitro* conditions, all concentrations of melaleuca essential oil reduced the mycelial growth of *A. alternata*. The oil reached complete inhibition of fungal growth from 0.2% concentration and above. In the cowpea treatment, the essential oil had no significant control over the percentage of infected seeds.

Conclusion: The melaleuca essential oil had a fungitoxic effect on the *A. alternata* under *in vitro* conditions. However, using the adopted methodology, on the cowpea bean seed treatment, the essential oil did not reduce the incidence of *A. alternata*.

Keywords: Alternative control; Cowpea bean; Mycelial growth; phytopathogenic fungi; tea-tree essential oil; seeds disease; *Vigna unguiculata*.

1. INTRODUCTION

Cowpea bean (*Vigna unguiculata* (L.) Walp), popularly known as the string bean or macaçar bean, is a source of protein and staple food for a large part of the population of the North and Northeast of Brazil, thus one of the most important crops in the country [1]. According to CONAB [2], Brazil occupies the third position in world bean production with a cultivation area of approximately one million hectares, with the North and Northeast regions accounting for about 90% of the cultivated area [3].

Cowpea cultivation has a very competitive production cost, a factor that has increased the farmers' interest in the crop. In addition, Brazilian production is of high quality, enabling the product to have good acceptance in all members of its production chain [4]. However, diseases represent a limiting factor to income expansion.

Fungi are the main phytopathogens that cause economic losses in bean crop. When present in the seed, they can cause miscarriages,

deformations and discoloration of the bark, which always leads to the reduction of seed germination potential and vigor, and when allocated in the field will result in low or no yielding uneven plant stands [5]. Diseases caused by fungi with the greatest economic impact on bean crop are caused by *Macrophomina phaseolina* [6], *Fusarium* spp. [7], *Rhizoctonia solani* [8], *Curvularia* spp., *Trichoderma* spp. [9], *Alternaria* spp. [10], *Aspergillus* sp. and *Penicillium* sp. [11].

Considering bean pathogen control practices, treatment with synthetic agrochemicals has been a conventionally used one. However, the use of these products has been associated with significant damage to human health and the environment due to their high toxicity [12,13] besides favoring the emergence of resistant strains [14].

In this scenario, it is necessary to use alternative products to chemical pesticides that have similar efficacy but are not harmful to human health and the environment. Among the products studied are essential oils extracted from aromatic plants,

which have fungitoxic properties on phytopathogens [15,16,17].

Melaleuca EO (*Melaleuca alternifolia*) has been studied for some years and its antimicrobial activity has been well documented. The main components of this oil are: terpinen-4-ol, cineol, terpenene, cymene, limonene and sabinene [18]. Most compounds have inhibitory activity against microorganisms [19], Terpinen-4-ol being the main constituent with antifungal activity [20]. In the control of phytopathogens its use has shown promising results in the control of fungi *Cercospora beticola* [21], *Aspergillus niger*, *M. phaseolina*, *Penicillium* sp. and *Sclerotinia sclerotiorum* [22], demonstrating a strong antimicrobial activity.

The antifungal action of essential oils is related to their ability to dissolve in lipid media, causing modifications in the cell membrane structure [23]. Due to their low toxicity and rapid degradation in the environment, the use of essential oils to combat phytopathogens may be a good alternative to synthetic pesticides [24]. Thus, this work aimed to evaluate the fungitoxic potential of melaleuca essential oil on the mycelial growth of *Alternaria alternata* under *in vitro* conditions and in the treatment of cowpea seeds.

2. MATERIALS AND METHODS

2.1 Place of Experiments

The work was conducted at the Center of Science and Technology Agrifood (CCTA) of the Federal University of Campina Grande (UFCG), Campus of Pombal. The experiments were carried out in the Phytopathology laboratory, between February 2018 and February 2019.

2.2 Sampling

The fungal strain used was *Alternaria alternata* 0878, which was provided by the collection of phytopathogenic fungi Prof. Maria Menezes of the Federal Rural University of Pernambuco (UFRPE). The fungi were preserved in sterile distilled water by the Castellani method until the assay [25].

The pure essential oil of Melaleuca (*Melaleuca alternifolia*) was purchased at a local store specialized in natural products. The cowpea bean seeds (*Vigna unguiculata* L. Walp) were purchased at a commercial house in the city of Patos, Paraíba.

2.3 Screening of the Antifungal Activity of Melaleuca Essential Oil *In vitro*

Eleven treatments were used, 9 oil concentrations (0.0125, 0.025, 0.1, 0.2, 0.5, 0.25, 0.50, 0.75 and 1.0%), a negative control (without essential oil supplementation=0.0%) and a positive control (supplemented with 1 mL L⁻¹ of the fungicide Thiram, which is the dosage indicated by the manufacture's). Five replicates of each treatment were arranged in completely randomized design (CRD).

The treatments were incorporated into PDA (Potato Dextrose Agar) culture medium just before pouring in sterilized Petri dishes. After solidification, one-centimeter mycelial disks were taken from the margins of 7 days old culture and transferred to the center of each plate containing the treatments. The plates were then wrapped in plastic film and incubated in a BOD (Biochemical Oxygen Demand) at a temperature of 27±2°C.

The concentrations were chosen from an initial concentration based on the literature [26,27] and then gradually reduced until the addition of oil to the medium was no longer able to prevent the fungal growth. To obtain the final concentrations, the direct dilution procedure in a culture medium [28] was used.

Colony growth was measured daily until the colony took the entire surface of the culture medium in one of the plates or in a maximum period of 7 days. Mycelial growth evaluation consisted of daily measurements of the diameter of the colonies obtained through the average of two perpendicular measurements, using a digital caliper, resulting in the average daily growth for each repetition of each treatment.

The percentage of mycelial growth inhibition (PGI; [29]) and mycelial growth rate index (IMGS; [30]) were calculated according to formulas (1) and (2):

$$PGI = \frac{[(negative\ control\ growth - treatment\ growth)] \times 100}{negative\ control\ growth} \quad (1)$$

$$IMGS = \frac{\sum (current\ mycelial\ growth - previous\ mycelial\ growth)}{number\ of\ days\ of\ incubation} \quad (2)$$

The minimum inhibitory concentration was considered the lowest oil concentration capable of totally inhibiting *Alternaria alternata* mycelial growth.

2.4 Screening of the Antifungal Activity of Melaleuca Essential Oil *In vivo* (on Cowpea Bean Seeds)

The experiment consisted of a completely randomized design. The treatments consisted of sterilized distilled water solutions supplemented with 4 oil concentrations (0.2, 0.5, 1.0 and 5.0%), a negative control (without essential oil supplementation=0.0%) and a positive control (supplemented with 1 ml L⁻¹ of the fungicide Thiram, which is the dosage indicated by the manufacturer's). The concentrations used were determined based on the *in vitro* test. To emulsify the oil in the water Tween 80 (1 mL L⁻¹) was used [31].

The cowpea bean seeds were disinfected in 2.0% sodium hypochlorite solution for five minutes, washed with sterile distilled water twice and dried at room temperature. Afterwards they were immersed for five minutes in different solutions (treatments). After drying at room temperature, the artificial inoculation was performed.

The inoculation was done depositing the seeds on 7 days colonies of *A. alternata*. The seeds and the fungal colonies stayed for 48 hours in a BOD 27±2°C, with a 12-hour photoperiod [32].

After the treatment and inoculation, the samples were submitted to the sanity test, which was performed by the filter paper method with freezing [33]. Six hundred of cowpea bean seeds (100 per treatment) were used, distributed in Petri dishes (Ø=14 cm). In this method, ten seeds were placed at equal distances on each plate on triple layer of filter paper previously moistened in sterile distilled water and incubated initially for 24 hours on BOD at 27±2°C, with a 12-hour photoperiod. After this period, they were subjected to freezing (-20°C) for 24 hours, and then returned to the incubator for another five days.

After incubation, the seed were examined individually, using a stereoscopic microscope, for the quantification of seeds infected by *Alternaria alternata*. The results were expressed as percentage of infected seeds.

2.5 Statistical Analysis

The effect of oil concentration on fungal growth was analyzed using regressions in quadratic

plateau model for *in vitro* experiment and in linear model for *in vivo* experiment.

To test the difference between treatments with the essential oil and the treatment containing the fungicide (positive control), Mann-Whitney (Tukey nonparametric) multiple comparisons were applied. Non-parametric tests were used because of the lack of variance in the results of some treatments. Differences with a probability values below 5% were considered significant. The regressions were performed in the program R Core Team 3.5.1 [34].

3. RESULTS AND DISCUSSION

3.1 *In vitro* Antifungal Assay

3.1.1 Effects of melaleuca essential oil on *Alternaria alternata*

All tested concentrations of melaleuca essential oil reduced the mycelial growth of *Alternaria alternata* (Fig. 1). The inhibition percentages increased significantly with the concentrations ($P<.001$) reaching the maximum value (PGI=100%) the 0.2% concentration of the oil (Fig. 2A), which is the minimum inhibitory concentration (MIC). On the other hand, applying the regression equation in a quadratic plateau model, the estimated minimum concentration (MCest) was 0.33%.

The mycelial growth rate is a variable inversely proportional to the inhibition percentage. For this reason, it presented opposite behavior, with significant reduction with the tested oil concentrations ($P<.001$). The mycelial growth rate was more effectively reduced from the 0.2% concentration, in which growth paralyzed (IMGS=0.00 cm day⁻¹) (Fig. 2B), differing from the negative control, which presented the highest growth speed (0.63 cm day⁻¹).

According to the literature, terpinen-4-ol is the major constituent of the melaleuca essential oil, which is associated with your high fungitoxic potential [35]. One of the antifungal mechanisms of action of melaleuca essential oil is the change in the permeability and fluidity of the cell membranes of the microorganisms. As these organisms are permeable to oil, the main effects found are inhibition of cell respiration and alteration in membrane structure and integrity, as well as leakage of essential intracellular materials. These events cause growth inhibition or even cell death [36,37].




Fig. 1. Comparison of *Alternaria alternata* mycelial growth in different treatments (melaleuca essential oil and the control treatments)

*A*¹ and *A*²: Negative control (without essential oil supplementation) and positive control (1 mL L⁻¹ of Thiram); *B*¹, *B*², *B*³, *B*⁴, *B*⁵, *B*⁶, *B*⁷, *B*⁸ and *B*⁹: Melaleuca essential oil at 0.0125, 0.025, 0.05, 0.1, 0.2, 0.25, 0.5, 0.75 and 1.0%, respectively




Fig. 2. Inhibition percentage and mycelial growth speed of melaleuca essential oil against *Alternaria alternata*

***Level of significance below 0.1% ($P < .001$)

Using tea tree oil at concentrations close to or greater than ours, other authors obtained similar inhibition results. For example, Martins et al. [22] obtained total inhibition of *M. phaseolina*

and *S. sclerotiorum* at concentration 0.2%. While in the control of *Alternaria radicina* and *A. dauci*, Rlcioni and Orzali [38] reached the maximum inhibition from the concentration 0.5%.

Using the essential oil of the other plant species on control of *A. alternata*, other authors obtained similar results as superior or inferior to ours. For example, the total inhibition was achieved by Chutia et al. [39], Guimarães et al. [40] and Barboza [41] using mandarin orange (*Citrus reticulata*), lemongrass (*Cymbopogon citratus*) and alecrim-da-chapada (*Lippia gracilis*) essential oil at concentrations of 0.2 mL/100mL (0.2%), 14.49 µg mL⁻¹ (0.0014%) e 750 µL L⁻¹ (0.075%), respectively. On the other hand, using peppermint essential oil (*Mentha piperita*), França et al. [42] obtained a maximum inhibition of 41.6% at a concentration of 0.8%. Thus, both the fungitoxic potential of essential oils on *A. alternata*, as well as their minimum inhibitory concentrations will vary depending on the plant species from which the essential oil was extracted [43]. In addition, increasing inhibitory power as a function of increased concentration can either potentiate the effect or generate product waste.

To understand the potential of melaleuca essential oil as a fungicide on *A. alternata*, the fungitoxic effects of the essential oil and the

fungicide Thiram (commercial synthetic fungicide) were compared. The essential oil, from the concentration of 0.2%, and the fungicide had similar inhibitions (Fig. 3). This result suggests that under *in vitro* conditions the oil could replace the use of this agrochemical.

Due to the chemical complexity, the antifungal control promoted by essential oils is associated with their different constituents [44] through different mechanisms of action that act simultaneously on different targets [15]. These peculiar characteristics guarantee the advantage over synthetic fungicide, since they reduce the possibility of resistance by phytopathogens [45].

3.2 In vivo Antifungal Assay

3.2.1 Effects of melaleuca essential oil on cowpea beans seed infected with *Alternaria alternata*

Contrary to the *in vitro* test, the melaleuca essential oil was impotent against the incidence of *A. alternata* in cowpea seeds ($P < .001$; Fig. 4). The incidence of infected seeds at concentrations from 0.2 to 1.0% was similar to the negative control ($P > .05$), while at 5% concentration the incidence of phytopathogen increased substantially (Fig. 5). On the other hand, the Thiram fungicide prevented the development of phytopathogen in the seeds.




Fig. 3. Effect of different treatments (melaleuca essential oil and the control treatments) on the mycelial growth inhibition of *Alternaria alternata*
 Superscript concentrations with the same letter were not significantly different from each other by the MannWhitney test ($P > .05$)




Fig. 4. Effect of concentrations of melaleuca essential oil in the incidence of infected cowpea bean seeds by *Alternaria alternata*




Fig. 5. Percentage of infected cowpea bean seeds by *Alternaria alternata* after the treatment with different concentrations of melaleuca essential oil and the control treatments

The potentiation of fungal growth in seeds treated with the melaleuca essential oil at 5% concentration may have occurred because of your low adherence to the surface of the seeds due to the high volatility of its constituents. Thus, during the incubation period some constituents may have evaporated and reduced to their antimicrobial capacity. The volatilization of oil constituents as well as their instability in the presence of light, heat and humidity, modify the atmosphere inside the Petri dishes, leading to the loss of the effectiveness of an oil that, under other conditions, inhibited fungal growth [46,47].

Another hypothesis would be that some innocuous constituents present in higher

concentrations became nutritive resources to the fungus, favoring its development even in the presence of fungitoxic components. Or, the high concentration of the essential oil may have affected the surface of the seeds and facilitated their colonization by the pathogen.

Despite our results, the use of other essential oil against phytopathogens in the bean seeds was successful. For example, the incidence of *Aspergillus* sp. and *Penicillium* sp. in beans treated with lemongrass (*Cymbopogon flexuosus* and *C. citratus*) and melaleuca (*Melaleuca* sp.) was reduced [48]. Also, citronella (*Cymbopogon* sp.), anise (*Pimpinella anisum*), and basil (*Ocimum basilicum*) essential oils at 1.5% concentration inhibited *Callosobruchus maculatus* [49].

Finally, despite the ineffectiveness of melaleuca essential oil in the treatment of cowpea bean seed, this oil may be useful in the treatment of other seeds and other pathogens. Essential oils present a low risk to the environment, producers and consumers, and hinder the development of pathogen resistance [50]. Thus, further studies on the use of these oils in the management of plant pathogens are needed to make them a viable alternative for farmers.

4. CONCLUSIONS

Under *in vitro* conditions, melaleuca essential oil (*Melaleuca alternifolia*) totally inhibited the mycelial growth of *Alternaria alternata* from 0.2%, had a similar effect to the commercial fungicide Thiram. On the cowpea bean seed treatment, the essential oil had was not able to reduce the

incidence of *A. alternata* using the adopted methodology.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Neves AC, et al. Cultivo do feijão-caupi em sistema agrícola familiar. Teresina- PI: Embrapa Meio Norte. (Embrapa Meio Norte. Circular Técnica, 51). Portuguese. 2011;15.
2. CONAB - Companhia Nacional de Abastecimento. Perspectiva para a Agropecuária. Brasília. 2018;6:1-112. Portuguese.
3. Santos LAC, Silva DMP, Oliveira IA, Pereira CE, Campos MCC. Crescimento de cultivares de feijão-caupi em solo de terra firme e várzea. *Ambiência Guarapuava*. 2017;13(1):261-270. Portuguese.
4. Freire Filho FR. Feijão-caupi no Brasil: Produção, melhoramento genético, avanços e desafios. Embrapa Meio-Norte, Teresina. 2011;84. Portuguese.
5. Machado JC. Tratamento de sementes no controle de doenças. Lavras: LAPS/UFLA/FAEPE. 2000;13.
6. Athayde Sobrinho C, Viana FMP, Santos A. Doenças fúngicas e bacterianas. In: Freire Filho FR, Lima JAA, Ribeiro VQ. Feijão-caupi: Avanços tecnológicos. (EDIÇÃO) Brasília, DF: Embrapa Informação Tecnológica, Capítulo. 2005; 12:461-484.
7. Rodrigues AAC, Menezes M. Detecção de fungos endofíticos em sementes de caupi provenientes de Serra Talhada e de Caruaru, estado de Pernambuco. *Fitopatologia Brasileira*. 2002;27(5):532-537. Portuguese.
8. Nechet KL, Halfeld-Vieira BA. Caracterização de isolados de *Rhizoctonia* spp., associados à mela do Feijão-Caupi (*Vigna unguiculata*), coletados em Roraima. *Fitopatologia Brasileira*. 2006;31: 505-508.
9. Sinha A, Singh SK, Qaisar J. Seed mycoflora of French bean and its control by means of fungicides. *Tropenlandwirt*. Witzenhousen. 1999;11(1):59-67.
10. Marques RO, Alves VM, Lima MLP, Arruda GMT. Avaliação sanitária e fisiológica de feijão oriundos de Unai – MG, Paracatu-MG e Cristalina-GO. *Summa Phytopathologica*. 2006;32(Suplemento). Portuguese.
11. Torres SB, Bringel JMM. Avaliação da qualidade sanitária e fisiológica de sementes de feijão macassar. *Caatinga*. 2005;18(2):88-92. Portuguese.
12. Candido e Silva EK, Melo LGL. Manejo de doenças de plantas: Um enfoque agroecológico. *Revista EDUCAmazônia–Educação Sociedade e Meio Ambiente*. 2013;10(1):143-157. Portuguese.
13. Perina FJ. Óleos essenciais e frações majoritárias ativas no controle da mancha marrom de alternaria (*Alternaria alternata*) em tangerina ponkan. 2014. 112f. Tese (Doutorado em Agronomia). Universidade Federal de Lavras, Lavras, MG; 2014. Portuguese.
14. Hillen T, Schwan-Estrada KRF, Mesquini RM, Cruz MES, Stangarlin JR, Nozaki M. Atividade antimicrobiana de óleos essenciais no controle de alguns fitopatógenos fúngicos *in vitro* e no tratamento de sementes. *Revista Brasileira de Plantas Mediciniais*. 2012;14(3):439-445. Portuguese.
15. AbdeL-Kader MM, El-Mougy NS, Lashin SM. Essential oils and *Trichoderma harzianum* as an integrated control measure against faba bean root rot pathogens. *Journal of Plant Protection Research*. 2011;51(3):306-311.
16. Silva M, Souza HRT, David HMSS, Santos LM, Silva RF, Amaro HTR. Qualidade fisiológica e armazenamento de sementes de feijão-comum produzidas no norte de Minas Gerais. *Revista Agro Ambiente*. 2014;8:97-103. Portuguese.
17. Santos MM, Peixoto AR, Pessoa ES, Nepa HBS, Paz CD; Souza AVV. Estudos dos constituintes químicos e atividade antibacteriana do óleo essencial de *Lippia gracilis* a *Xanthomonas campestris* pv. viticola “*in vitro*”. *Summa Phytopathologica*, Botucatu. 2014;40(3): 277-280. Portuguese.
18. Carson CF, Hammer KA, Riley TV. *Melaleuca alternifolia* (tea tree) oil: A review of antimicrobial and other medicinal properties. *Clinical Microbiology Reviews*. 2006;19(1):50-62.
19. Sagave L, Gressler LT, Flores FC, Silva CB, Vargas APC, Lovato M, Sangioni LA, Pötter L, Botton SA. Atividade de nanoformulações de *Melaleuca alternifolia*

- e terpinen-4-ol em isolados de *Rhodococcus equi*. Arquivo Brasileiro de Medicina Veterinária e Zootecnia. 2015; 67(1):221-226. Portuguese.
20. Vieira TR, Barbosa LCA, Maltha CRA, Paula VF, Nascimento EA. Constituintes químicos de *Melaleuca alternifolia* (Myrtaceae). Química Nova. 2004;27(4): 536-539. Portuguese.
 21. Souza AD, Roggerio TU, Furlan MR, Aoyama EM. Óleo de melaleuca (*Melaleuca alternifolia* Maiden & Betche, Cheel) no controle de cercosporiose em beterraba. Revista Brasileira de Plantas Mediciniais. 2015;17(4):1078-1082. Portuguese.
 22. Martins JAS, Sagata E, Santos VA, Juliantti FC. Avaliação do efeito do óleo de *Melaleuca alternifolia* sobre o crescimento micelial *in vitro* de fungos fitopatogênicos. Bioscience Journal. 2010;27(1):49-51a. Portuguese.
 23. Brum RBCS, Cardon CH, Pereira AS, Cardoso DP, Dos Santos GR. Atividade antifúngica de óleos essenciais sobre fungos fitopatogênicos. Magistra. 2014; 26(3):361-371. Portuguese.
 24. Silva BVS, Paiva GF, Barbieri TPOM, Souza GHS, Gonçalves FJT. Efeito dos Óleos Essenciais Sobre o Crescimento Micelial *in vitro* de *Colletotrichum* spp. Cadernos de Agroecologia. 2018;13(2): 1-9. Portuguese.
 25. Castellani A. Maintenance and cultivation of common pathogenic fungi of man in sterile distilled water. Further Researches. Journal of Tropical Medicine and Hygiene. 1967;70:181-84.
 26. Concha JM, Moore LS, Holloway WJ. Antifungal activity of *Melaleuca alternifolia* (tea-tree) oil against various pathogenic organisms. Journal. American Podiatric Medical Association. 1998;88(10):489-92.
 27. Martins ESCS, Santos MS, Barros HMM, Farias MAA. Efeito dos óleos essenciais de citronela, alecrim e erva-cidreira no controle *in vitro* de *Ralstonia solanacearum* em pimentão. Tecnologia & Ciência Agropecuária. 2010;4(1):9-13b. Portuguese.
 28. Pereira MC, Vilela GR, Costa LMAS, Silva RF, Fernandes AF, Fonseca EWNF, Picolli RH. Inibição do desenvolvimento fúngico através da utilização de óleos essenciais de condimentos. Ciência e Agrotecnologia. 2006;30(4):731-738. Portuguese.
 29. Bastos CN. Efeito do óleo de *Piper aduncum* sobre *Crinipelis pernicioso* e outros fungos fitopatogênicos. Fitopatologia Brasileira. 1997;22(3):441-443. Portuguese.
 30. Oliveira JA. Efeito do tratamento fungicida em sementes no controle de tombamento de plântulas de pepino (*Cucumis sativas* L.) e pimentão (*Capsicum annum* L.). 1991. 111 f. Dissertação (Mestrado em Fitossanidade) – Escola Superior de Agricultura de Lavras, Lavras; 1991. Portuguese.
 31. Santos PL. Manejo de *Macrophomina phaseolina* (Tassi) Goid. em sementes de feijoeiro (*Phaseolus vulgaris* L.) com óleos essenciais e antagonistas. 76fls. Tese (Doutorado em Agronomia) – Universidade Estadual Paulista, Botucatu-SP; 2018. Portuguese.
 32. Silva MAD, Silva WR. Comportamento de fungos e de sementes de feijoeiro durante o teste de envelhecimento artificial. Pesquisa Agropecuária Brasileira. 2000; 35(3):599-608. Portuguese.
 33. Limonard T. A modified blotter test for seed health. Netherland Journal of Plant Pathology. 1996;72(2):319-321.
 34. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2018. Available:<https://www.R-project.org/>
 35. Hammer KA, Carson CF, Riley TV. Effects of *Melaleuca alternifolia* (Tea Tree) essential oil and the major monoterpene component Terpinen-4-ol on the development of single and multistep antibiotic resistance and antimicrobial susceptibility. Antimicrobial Agents and Chemotherapy. 2012;52(2):909-915.
 36. Hammer KA, Carson CF, Riley TV., Antifungal effects of *Melaleuca alternifolia* (Tea tree) oil and its components on *Candida albicans*, *Candida glabrata* and *Saccharomyces cerevisiae*. Journal of Antimicrobial Chemotherapy. 2004;53: 1081-1085.
 37. Azevedo FA, Polydoro DA, Bastianel M, Kupper KC, Stuart RM, Costa FP, Pio RM. Resposta de diferentes genótipos de tangerinas e seus híbridos à inoculação *in vitro* e *in vivo* de *Alternaria alternata*. Revista Brasileira de Fruticultura. 2010; 32:1-8. Portuguese.
 38. Riccioni L, Orzali L. Activity of tea tree (*Melaleuca alternifolia*, Cheel) and thyme

- (*Thymus vulgaris*, Linnaeus.) essential oils against some pathogenic seed borne fungi. Journal of Essential Oil Research. 2011; 23(6):43-47.
39. Chutia M, Bhuyan PD, Pathak MG, Sharma TC, Boruah P. Antifungal activity and chemical composition of *Citrus reticulata* Blanco essential oil against phytopathogens from North East India. LWT - Food Science and Technology. 2009;42:777-780.
 40. Guimarães LGL, Cardoso MG, Sousa PE, Andrade J, Vieira SS. Atividades antioxidante e fungitóxica do óleo essencial de capim-limão e do citral. Revista Ciência Agronômica. 2011; 42(2):464-472. Portuguese.
 41. Barboza HS. Efeito fungitóxico do óleo essencial de alecrim-da-chapada em *Alternaria* sp. 2015. 31fls. Trabalho de Conclusão de Curso (Graduação em Agronomia) – Universidade Federal do SemiÁrido, Mossoró, RN, Brasil; 2015. Portuguese.
 42. França KRS, Lima TS, Cardoso TAL, Ugulino ALN, Rodrigues APMS, Mendonça Júnior AF. *In vitro* Effect of essential oil of peppermint (*Mentha x piperita* L.) on the mycelial growth of *Alternaria alternata*. Journal of Experimental Agriculture International. 2018;26(5):1-7.
 43. Antunes MDC, Cavacob A. The use of essential oils for postharvest decay control. A review. Flavour Fragrance Journal. 2010;25:351-366.
 44. Russo M, Suraci F, Postorino S, Serra D, Roccotelli A, Agosteo GE. Essential oil chemical composition and antifungal effects on *Sclerotium cepivorum* of *Thymus capitatus* wild populations from Calabria, southern Italy. Revista Brasileira de Farmacognosia. 2013;23(2): 239-248.
 45. Feng W, Zheng X. Essential oils to control *Alternaria alternata* *in vitro* and *in vivo*. Food Control. 2007;18:1126-1130.
 46. Simões CMO, Spitzer V. Óleos voláteis. In: Gosmann G, Mello JCP, Simões CMO, Schenkel EP. (Org.). Farmacognosia: da planta ao medicamento. 5th ed. Porto Alegre: UFRGS/UFSC. 2004;467-495. Portuguese.
 47. Rozwalka LC. Extratos, decotos e óleos essenciais de plantas medicinais e aromáticas na inibição de *Glomerella cingulata* e *Colletotrichum gloeosporioides* de frutos de goiaba. Ciência Rural. 2008;58(2):301-307. Portuguese.
 48. Morais LAS, Ramos NP, Gonçalves GG, Betiol W, Chaves FCM. Atividade antifúngica dos óleos essenciais em sementes de feijão cv. Carioquinha. Horticultura Brasileira. 2008;26(2):6261-6266. Portuguese.
 49. Wanderley MJA, Silva TMB, Arruda JD, Rocha RB, Cruz CSA, Wanderley PA. Efeito de óleos essenciais sobre *Callosobruchus maculatus* em feijão caupi *Vigna unguiculata* L. Walp. In: 6º Congresso Brasileiro de Plantas Oleaginosas, Óleos, Gorduras e Biodiesel. 2009. Minas Gerais, Montes Claros; 2009. Portuguese.
 50. Derbahlah AS, Hhamza AM, Gazzy AA. Efficacy and safety of some plant extracts as alternatives for *Sitophilus oryzae* control in rice grains. Journal of Entomology. 2012;9:57-67.

© 2019 Alves et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50927>